

A stability transfer theorem in d-Tame Metric Abstract Elementary Classes

Pedro Zambrano

Universidad Nacional de Colombia
Bogota - Colombia


XV SLALM, Bogota, Colombia
June 4th, 2012

- 1 Motivation
- 2 Definition of MAEC
- 3 d -Tameness and independence in MAEC
- 4 Stability transfer theorems

Discrete *tame* AEC are a special kind of AEC which have a categoricity transfer theorem (due to Grossberg and VanDieren) and a nice stability transfer theorem (due to Baldwin, Kueker and VanDieren), by using ω -locality.

MAEC corresponds to a kind of amalgam between AEC and *Continuous Logic Elementary Classes*. In this work, we study a version of tameness in this setting and prove a stability transfer theorem, removing the ω -locality assumption and assuming local character of a suitable well-behaved notion of independence.

A map of discrete and metric structures classes



Definition

Let \mathcal{K} be a class of L -structures (in the setting of Continuous Logic, but the function symbols need not be uniformly continuous). and let $<_{\mathcal{K}}$ be a binary relation defined on \mathcal{K} . We say that $(\mathcal{K}, <_{\mathcal{K}})$ is a *Metric Abstract Elementary Class* (for short, *MAEC*) iff:

- (1) \mathcal{K} and $<_{\mathcal{K}}$ are closed under \cong
- (2) $<_{\mathcal{K}}$ is a partial order in \mathcal{K} .
- (3) If $\mathcal{M} <_{\mathcal{K}} \mathcal{N}$ then $\mathcal{M} \subseteq \mathcal{N}$.

Definition (MAEC)

- (4) (Tarski-Vaught chains) If $(\mathcal{M}_i : i < \lambda)$ is an increasing and continuous $<_{\mathcal{K}}$ -chain, then
- the function symbols in L can be uniquely interpreted in the completion of $\bigcup_{i < \lambda} \mathcal{M}_i$ such that $\overline{\bigcup_{i < \lambda} \mathcal{M}_i} \in \mathcal{K}$
 - for all $j < \lambda$, $\mathcal{M}_j <_{\mathcal{K}} \overline{\bigcup_{i < \lambda} \mathcal{M}_i}$
 - if for every $\mathcal{M}_i \in \mathcal{K} <_{\mathcal{K}} \mathcal{N}$, then $\overline{\bigcup_{i < \lambda} \mathcal{M}_i} <_{\mathcal{K}} \mathcal{N}$.
- (5) (coherence) If $\mathcal{M}_1 \subseteq \mathcal{M}_2 <_{\mathcal{K}} \mathcal{M}_3$ and $\mathcal{M}_1 <_{\mathcal{K}} \mathcal{M}_3$, then $\mathcal{M}_1 <_{\mathcal{K}} \mathcal{M}_2$.
- (6) (DLS) There is a cardinality $LS(K)$ (which is called *Löwenheim-Skolem number of \mathcal{K}*) such that if $\mathcal{M} \in \mathcal{K}$ and $A \subseteq M$, then there exists $\mathcal{N} \in \mathcal{K}$ such that $dc(\mathcal{N}) \leq dc(A) + LS(K)$ and $A \subseteq \mathcal{N} <_{\mathcal{K}} \mathcal{M}$.

Examples of MAECs

- 1 The class of Banach spaces
- 2 The subclass of complete models in an elementary class of a positive bounded theory (W. Henson - J. Iovino).
- 3 Compact Abstract Theories (I. Ben-Yaacov).
- 4 $(Mod(T), <)$, T a (first order) theory in Continuous Logic:
 - 1 Hilbert spaces with unitary operators (C. Argoty - A. Berenstein)
 - 2 Nakano spaces with compact essential rank (P. Poitevin).
- 5 AECs (with the discrete metric)

Remark

Under AP+JEP+existence of large enough models, we can work in a homogeneous monster model $\mathbb{M} \in \mathcal{K}$.

Definition

$ga - tp(a/M)$ is defined as the orbit of the element a under automorphisms of \mathbb{M} which fix M pointwise.

Definition

$ga-S(M) := \{ga-tp(a/M) : a \in \mathbb{M}\}$

Some preliminary results

Definition

Let $p, q \in \text{ga-S}(M)$ ($M \in \mathcal{K}$). $d(p, q) := \inf\{d(a, b) : a \models p, b \models q\}$.

Definition

We say that a MAEC satisfies the *continuity* property (for short, *CP*) iff $(a_n) \rightarrow b$ and $\text{ga-tp}(a_0/M) = \text{ga-tp}(a_n/M)$ for every $n < \omega$ implies that $\text{ga-tp}(b/M) = \text{ga-tp}(a_0/M)$.

Fact (Hirvonen-Hyttinen)

d is a metric in $\text{ga-S}(M)$ ($M \in \mathcal{K}$) iff \mathcal{K} satisfies the *CP*.

Definition

We say that a MAEC \mathcal{K} is μ -**d**-stable iff for every $M \in \mathcal{K}$ of density character μ we have that $dc(\text{ga-S}(M)) \leq \mu$.

Cofinal d-stability

Let \mathcal{K} be an MAEC with AP and JEP and $LS(\mathcal{K}) \leq \lambda < \kappa$. We say that \mathcal{K} is $[\lambda, \kappa)$ -*cofinally d-stable* iff given $\theta \in [\lambda, \kappa)$ there exists $\theta' \geq \theta$ in $[\lambda, \kappa)$ such that \mathcal{K} is θ' -**d**-stable.

Tameness (discrete AEC)

Let \mathcal{K} be an AEC and $\mu \geq LS(\mathcal{K})$. We say that \mathcal{K} is μ -tame iff for any $M \in \mathcal{K}$ of cardinality $\geq \mu$, if $p \neq q$ where $p, q \in \text{ga-S}(M)$, then there exists $N <_{\mathcal{K}} M$ of cardinality μ such that $p \upharpoonright N \neq q \upharpoonright N$.

Tameness (discrete AEC)

Let \mathcal{K} be an AEC and $\mu \geq LS(\mathcal{K})$. We say that \mathcal{K} is μ -tame iff for any $M \in \mathcal{K}$ of cardinality $\geq \mu$, if $p \neq q$ where $p, q \in \text{ga-S}(M)$, then there exists $N \prec_{\mathcal{K}} M$ of cardinality μ such that $p \upharpoonright N \neq q \upharpoonright N$.

d-Tameness

Let \mathcal{K} be a MAEC and $\mu \geq LS(\mathcal{K})$. We say that \mathcal{K} is μ -**d**-tame iff for every $\varepsilon > 0$ there exists $\delta_\varepsilon > 0$ such that if for any $M \in \mathcal{K}$ of $\text{dc} \geq \mu$ we have that $\mathbf{d}(p, q) \geq \varepsilon$ where $p, q \in \text{ga-S}(M)$, then there exists $N \prec_{\mathcal{K}} M$ of $\text{dc} \mu$ such that $\mathbf{d}(p \upharpoonright N, q \upharpoonright N) \geq \delta_\varepsilon$.

Tameness (discrete AEC)

Let \mathcal{K} be an AEC and $\mu \geq LS(\mathcal{K})$. We say that \mathcal{K} is μ -tame iff for any $M \in \mathcal{K}$ of cardinality $\geq \mu$, if $p \neq q$ where $p, q \in \text{ga-S}(M)$, then there exists $N \prec_{\mathcal{K}} M$ of cardinality μ such that $p \upharpoonright N \neq q \upharpoonright N$.

d-Tameness

Let \mathcal{K} be a MAEC and $\mu \geq LS(\mathcal{K})$. We say that \mathcal{K} is μ -**d-tame** iff for every $\varepsilon > 0$ there exists $\delta_\varepsilon > 0$ such that if for any $M \in \mathcal{K}$ of $\text{dc} \geq \mu$ we have that $\mathbf{d}(p, q) \geq \varepsilon$ where $p, q \in \text{ga-S}(M)$, then there exists $N \prec_{\mathcal{K}} M$ of $\text{dc} \mu$ such that $\mathbf{d}(p \upharpoonright N, q \upharpoonright N) \geq \delta_\varepsilon$.

Some assumptions (*)

We assume \mathcal{K} is a μ -d-tame ($\mu < \kappa$) and $[LS(\mathcal{K}), \kappa)$ -cofinally-d-stable. Define $\lambda := \min\{\mu < \chi < \kappa : \mathcal{K} \text{ is } \chi\text{-d-stable}\}$ $\zeta := \min\{\chi : 2^\chi > \lambda\}$ and $\zeta^* := \max\{\mu^+, \zeta\}$. We also require that $\text{cf}(\kappa) \geq \zeta^*$ and $\kappa > \zeta^*$.

Definition (Splitting, AEC)

Let $N <_{\mathcal{K}} M$. We say that $\text{ga-tp}(a/M)$ splits over N iff there exist N_1 and N_2 such that $N <_{\mathcal{K}} N_1, N_2 <_{\mathcal{K}} M$ and $h : N_1 \cong_N N_2$ such that $\text{ga-tp}(a/N_2) \neq h(\text{ga-tp}(a/N_1))$.


Definition (Splitting, AEC)

Let $N <_{\mathcal{K}} M$. We say that $\text{ga-tp}(a/M)$ splits over N iff there exist N_1 and N_2 such that $N <_{\mathcal{K}} N_1, N_2 <_{\mathcal{K}} M$ and $h : N_1 \cong_N N_2$ such that $\text{ga-tp}(a/N_2) \neq h(\text{ga-tp}(a/N_1))$.

Definition (ε -splitting)

Let $N <_{\mathcal{K}} M$ and $\varepsilon > 0$. We say that $\text{ga-tp}(a/M)$ ζ^* - ε -splits over N iff for every $N^* <_{\mathcal{K}} N$ of $\text{dc} < \zeta^*$ there exist N_1 and N_2 of $\text{dc} < \zeta^*$ such that $N^* <_{\mathcal{K}} N_1, N_2 <_{\mathcal{K}} M$ and $h : N_1 \cong_{N^*} N_2$ such that $\mathbf{d}(\text{ga-tp}(a/N_2), h(\text{ga-tp}(a/N_1))) \geq \varepsilon$. If $\text{ga-tp}(a/M)$ does not tame- ε -split over N , we denote that by $a \downarrow_N^{\varepsilon} M$.

Independence in tame MAEC



Definition

Let $N <_{\mathcal{K}} M$. We say that a is ζ^* -independent from M over N iff for every $\epsilon > 0$ $a \downarrow_N^{\epsilon} M$.

Some properties of ε -independence

Some properties

- 1 (Local character I) For every M , a and every $\varepsilon > 0$ there exists $N \prec_{\mathcal{K}} M$ of density character $< \zeta^*$ such that $a \downarrow_N^{\varepsilon} M$.
- 2 (Weak stationarity) For every $\varepsilon > 0$ there exists δ such that for every $N_0 \prec_{\mathcal{K}} N_1 \prec_{\mathcal{K}} N_2$ and every a, b , if N_1 is universal over N_0 , $a, b \downarrow_{N_0}^{\delta} N_2$ and $\mathbf{d}(\text{ga-tp}(a/N_1), \text{ga-tp}(b/N_1)) < \delta$, therefore $\mathbf{d}(\text{ga-tp}(a/N_2), \text{ga-tp}(b/N_2)) < \varepsilon$.

More assumptions (**) -local character II-

For every tuple \bar{a} , every $\varepsilon > 0$ and every increasing and continuous $\prec_{\mathcal{K}}$ -chain of models $\langle M_i : i < \sigma \rangle$, there exists $j < \sigma$ such that $\bar{a} \downarrow_{M_j}^{\varepsilon} \overline{\bigcup_{i < \sigma} M_i}$.

Stability transfer theorems

Assumption (*)

We assume \mathcal{K} is μ -d-tame ($\mu < \kappa$) and $[LS(\mathcal{K}), \kappa)$ -cofinally-d-stable. Define $\lambda := \min\{\mu < \chi < \kappa : \mathcal{K} \text{ is } \chi\text{-d-stable}\}$ $\zeta := \min\{\chi : 2^\chi > \lambda\}$ and $\zeta^* := \max\{\mu^+, \zeta\}$. We also require that $cf(\kappa) \geq \zeta^*$ and $\kappa > \zeta^*$.

Theorem (#)

Let \mathcal{K} be an MAEC satisfying assumption (*). Then \mathcal{K} is κ -d-stable.

Idea of the proof.

By RA, using local character I, $cf(\kappa) \geq \zeta^*$ and pigeon-hole principle, contradicting cofinal stability. \square

Remark

If $\mu = \aleph_0$ and $\lambda = \aleph_1$, then $\zeta := \min\{\chi : 2^\chi > \lambda\} \leq \aleph_1$ and $\zeta^* = \aleph_0$.

Corollary

Let \mathcal{K} be an \aleph_0 -**d**-tame MAEC. Suppose that \mathcal{K} is \aleph_0 -*d*-stable and \aleph_1 -*d*-stable. Then \mathcal{K} is \aleph_n -*d*-stable for all $n < \omega$

Corollary (##)

Let \mathcal{K} be an \aleph_0 -**d**-tame MAEC. Suppose that \mathcal{K} is \aleph_0 -*d*-stable and \aleph_1 -*d*-stable. Then \mathcal{K} is \aleph_ω -*d*-stable.

Idea of the proof

By RA, use local character II, \aleph_0 -*d*-tameness and pigeon-hole principle to contradict \aleph_n -*d*-stability. □

Proposition

Let \mathcal{K} be an \aleph_0 -d-tame, \aleph_0 -d-stable and \aleph_1 -d-stable MAEC, which also satisfies assumption (**). Then \mathcal{K} is κ -d-stable for every cardinality κ .





Idea of the proof.

If $cf(\kappa) \geq \zeta^* = \omega_1$, use theorem \sharp . If $cf(\kappa) = \omega$, use a similar argument as in corollary ($\sharp \sharp$)


Acknowledgment

I thank Andrés Villaveces, Tapani Hyttinen and John Baldwin for the nice discussions and suggestions to improve this work.

References

-  Å. Hirvonen, T. Hyttinen, *Categoricity in homogeneous complete metric spaces*, Arch. Math. Logic 48, pp. 269–322, 2009.
-  J. Baldwin, D. Kueker, M. Vandieren, *Upward stability transfer theorem in tame AECs*, Notre Dame J. Formal Logic vol 47 n. 2, pp. 291–298, 2006.
-  P. Zambrano, *Around superstability in metric abstract elementary classes*, Ph.D. thesis (U. Nacional de Colombia), 2011.
-  P. Zambrano, *A stability transfer theorem in d-tame metric abstract elementary classes*, accepted at Math. Logic Quarterly, 2012

THANKS!



View from Colpatria tower - Bogotá.